

**Table 7.6d**  
**Immunosuppression Use for Maintenance by Regimen**  
**Prior to Discharge, 1998 to 2007**  
**Recipients with Pancreas After Kidney (PAK) Transplants**

	Transplant Year									
	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007
<b>Functioning Graft at Discharge</b>	139	206	277	287	336	314	380	305	256	240
<b>With Immunosuppression Info</b>	137	203	276	286	335	311	374	301	250	228
CyA	0.0%	0.0%	0.0%	0.3%	0.3%	0.0%	0.0%	0.0%	0.0%	0.0%
Tac	0.0%	6.9%	4.0%	3.8%	4.8%	1.3%	2.7%	3.0%	2.4%	2.6%
--- + MMF/MPA	0.0%	0.0%	0.0%	0.0%	0.0%	5.8%	5.1%	2.0%	0.0%	0.9%
CyA + MMF/MPA	0.0%	0.5%	0.0%	0.7%	0.3%	1.0%	1.3%	2.0%	0.8%	0.9%
Tac + MMF/MPA	0.0%	0.5%	0.0%	0.3%	0.9%	6.4%	8.0%	17.6%	23.2%	28.9%
Tac + OtherAntimet	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	0.0%	0.0%
--- + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.5%	0.3%	0.0%	0.0%
CyA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	1.1%	1.0%	2.4%	0.4%
Tac + Siro/Evero	0.0%	0.0%	0.0%	1.0%	2.4%	2.3%	4.0%	4.3%	2.4%	0.0%
--- + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	1.6%	0.3%	0.4%	0.4%
CyA + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.0%	0.3%	0.0%	0.0%
Tac + MMF/MPA + Siro/Evero	0.0%	0.0%	0.0%	0.0%	0.3%	1.6%	0.8%	1.7%	4.0%	7.0%
--- + Steroids	0.7%	2.0%	1.8%	0.3%	0.3%	1.0%	2.7%	3.3%	0.0%	0.0%
CyA + Steroids	1.5%	0.5%	0.0%	0.0%	0.3%	0.3%	0.5%	0.0%	0.4%	0.4%
Tac + Steroids	3.6%	2.5%	4.3%	1.0%	2.4%	1.3%	2.1%	0.3%	0.8%	1.8%
--- + MMF/MPA + Steroids	0.7%	1.0%	0.0%	0.3%	1.5%	1.6%	1.6%	0.3%	0.4%	1.3%
CyA + MMF/MPA + Steroids	18.2%	10.3%	10.9%	5.6%	7.2%	5.1%	3.5%	2.3%	2.0%	1.8%
Tac + MMF/MPA + Steroids	65.7%	68.5%	67.4%	62.6%	60.6%	59.8%	55.1%	53.8%	52.8%	51.3%
CyA + OtherAntimet + Steroids	0.7%	0.0%	0.0%	0.0%	0.0%	0.3%	0.0%	0.0%	0.0%	0.4%
Tac + OtherAntimet + Steroids	1.5%	2.0%	1.8%	0.7%	0.3%	0.3%	0.3%	0.7%	0.0%	0.4%
--- + Siro/Evero + Steroids	0.0%	0.0%	0.0%	0.0%	0.3%	0.3%	0.3%	0.3%	0.4%	0.0%
CyA + Siro/Evero + Steroids	0.7%	0.5%	1.8%	1.4%	2.1%	1.3%	0.8%	0.7%	0.0%	0.0%
Tac + Siro/Evero + Steroids	0.0%	1.0%	5.4%	15.7%	11.3%	6.1%	4.5%	2.7%	6.0%	0.0%
Other Regimen	6.6%	3.9%	2.5%	5.9%	4.8%	3.9%	3.5%	2.7%	1.6%	1.3%

Source: OPTN/SRTR Data as of May 1, 2008.

Notes:

Percentages in each column sum to 100 percent.

---: No Calcineurin Inhibitor Use; CyA: Cyclosporine; Tac: Tacrolimus.

MMF: Mycophenolate Mofetil; MPA: Mycophenolate Sodium; Siro: Sirolimus; Evero: Everolimus; OtherAntimet: Antimetabolites excluding MMF and MPA.

"Other Regimen" includes drug combinations that are different from the listed regimens and drugs from the text field entering.